

病名: 卵巣癌

レジメン名: CPT 100mg/m²

1コース期間(休薬期間含む) 28 日

薬剤名/ルート・用法	基準投与量	割合	1/1	1/2	1/3	1/4	1/5	1/6	1/7	1/8	1/9	1/10	1/11	1/12	1/13	1/14	1/15	1/16	1/17	1/18	1/19	1/20	1/21	1/22	1/23	1/24	1/25	1/26	1/27	1/28
			火	水	木	金	土	日	月	火	水	木	金	土	日	月	火	水	木	金	土	日	月	火	水	木	金	土	日	月
			day1 未適用	day2~day7 休薬					day8 未適用	day9~day14 休薬					day15 未適用	day16~day28 休薬														
注射																														
グラセトロン注 3mg100mL	1 V																													
デキサート注射液6.5mg 2mL	1 V																													
デキサート注 ★3.3mg 1mL	1 A																													
末梢点滴ケモ(持続)	00:05		未適用																											
生理食塩液注 500mL(テルパック)	1 V																													
●イリヌマクシ酸塩注 40mg 2mL	100 mg/m ²	100																												
末梢点滴ケモ CSTD(側管)	00:10		未適用																											
生理食塩液[ヒカリ]50mL	1 V																													
末梢点滴ケモ(持続)	00:15		未適用																											
グラセトロン注 3mg100mL	1 V																													
デキサート注射液6.5mg 2mL	1 V																													
デキサート注 ★3.3mg 1mL	1 A																													
末梢点滴ケモ(持続)	00:05									未適用																				
生理食塩液注 500mL(テルパック)	1 V																													
●イリヌマクシ酸塩注 40mg 2mL	100 mg/m ²	100																												
末梢点滴ケモ CSTD(側管)	00:10									未適用																				
生理食塩液[ヒカリ]50mL	1 V																													
末梢点滴ケモ(持続)	00:15									未適用																				
グラセトロン注 3mg100mL	1 V																													
デキサート注射液6.5mg 2mL	1 V																													
デキサート注 ★3.3mg 1mL	1 A																													
末梢点滴ケモ(持続)	00:05																													未適用
生理食塩液注 500mL(テルパック)	1 V																													
●イリヌマクシ酸塩注 40mg 2mL	100 mg/m ²	100																												
末梢点滴ケモ CSTD(側管)	00:10																													未適用
生理食塩液[ヒカリ]50mL	1 V																													
末梢点滴ケモ(持続)	00:15																													未適用

注意コメント
 day2-4 テ`カ`ロン 8mg/day