

病名: 胃癌

レジメン名: XELOX+トラスツズマブ(2回目以降)

1コース期間(休薬期間含む) 21 日

薬剤名/ルート・用法	基準投与量	割合	1/1	1/2	1/3	1/4	1/5	1/6	1/7	1/8	1/9	1/10	1/11	1/12	1/13	1/14	1/15	1/16	1/17	1/18	1/19	1/20	1/21
			火	水	木	金	土	日	月	火	水	木	金	土	日	月	火	水	木	金	土	日	月
			day1 未適用	day2~day21 休薬																			
注射																							
大塚生食注 ☆250mL	1 袋																						
トラスツズマブES点滴静注 ☆ 60mg	6 mg/kg	100																					
注射用水 20mL	20 ml																						
末梢点滴ケモ(持続)	00:10		未適用																				
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴(持続)	00:15		未適用																				
ハロピドロン点滴静注バッグ 0.75mg 50ml	1 袋																						
デキサート注 ★3.3mg 1mL	1 A																						
アロカリス点滴静注235mg10mL	1 V																						
末梢点滴ケモ(持続)加算A	00:20		未適用																				
ブドウ糖注 5%★500mL	1 V																						
オキサリプラチン注 ☆50mg	130 mg/m ²	100																					
デキサート注 ☆1.65mg 0.5mL	1 A																						
末梢点滴ケモ CSTD(側管)	00:25		未適用																				
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴ケモ(持続)	00:30		未適用																				
処方																							
注意コメント	day1(夕)~15(朝) カペシタビン2000mg/m ² /day 朝・夕、day2-4デキサロン4mg朝																						