

病名:前立腺癌

レジメン名:低用量ED(エストラムスチン・ドセタキセル)療法

1コース期間(休薬期間含む)

21 日

薬剤名/ルート・用法	基準投与量	割合	1/1	1/2	1/3	1/4	1/5	1/6	1/7	1/8	1/9	1/10	1/11	1/12	1/13	1/14	1/15	1/16	1/17	1/18	1/19	1/20	1/21		
			火	水	木	金	土	日	月	火	水	木	金	土	日	月	火	水	木	金	土	日	月		
			day1	day2	day3~day8					day9	day10~day21														
			休薬	未適用	休薬					未適用	休薬														
注射																									
生理食塩液「ヒカリ」 100mL	1 V																								
デキサート注射液6.6mg 2mL	1 V																								
ファミチジン注射液 20mg	1 A																								
末梢点滴ケモ(持続)	00:05			未適用																					
ブドウ糖注 5% 250mL	1 V																								
●ドセタキセル注 ☆20mg	25 mg/m ²	100																							
末梢点滴ケモ CSTD(側管)	00:10			未適用																					
生理食塩液「ヒカリ」 50mL	1 V																								
末梢点滴ケモ(持続)	00:15			未適用																					
生理食塩液「ヒカリ」 100mL	1 V																								
デキサート注射液6.6mg 2mL	1 V																								
ファミチジン注射液 20mg	1 A																								
末梢点滴ケモ(持続)	00:05										未適用														
ブドウ糖注 5% 250mL	1 V																								
●ドセタキセル注 ☆20mg	25 mg/m ²	100																							
末梢点滴ケモ CSTD(側管)	00:10										未適用														
生理食塩液「ヒカリ」 50mL	1 V																								
末梢点滴ケモ(持続)	00:15										未適用														
処方																									
注意コメント	day1-3、day8-10 朝夕 エストラサイト 4cap day1-21 朝 デカロン0.5mg 2T																								